

AD A13H 390

NEGATIVE-ION FORMATION FROM SURFACE SCATTERING AND THE  
ANDERSON CORRELATION (U) ROCHESTER UNIV NY DEPT OF  
CHEMISTRY K S LAM ET AL. FEB 84 UROCHESTER/DC/84/TR 49  
N00014-80-C-0472

1/1

UNCLASSIFIED

F/G 7/4

NI


END

DATE

FILED

3 84

DT



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT No. 49

Negative-Ion Formation from Surface Scattering  
and the Anderson Correlation Energy U

by

Kai-Shue Lam, K. C. Liu and Thomas F. George

Prepared for Publication

in

Physics Letters A

Department of Chemistry  
University of Rochester  
Rochester, New York 14627

February 1984

Reproduction in whole or in part is permitted for any  
purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

DTIC  
ELECTE  
FEB 24 1984  
S E D

DTIC FILE COPY

84 02 23 005

ADA138390

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UROCHESTER/DC/84/TR-49	2. GOVT ACCESSION NO. AD-A138390	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Negative-Ion Formation from Surface Scattering and the Anderson Correlation Energy U		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Kai-Shue Lam, K. C. Liu and <u>Thomas F. George</u>		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Rochester Rochester, New York 14627		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		12. REPORT DATE February 1984
		13. NUMBER OF PAGES 17
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication Physics Letters A		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) NEGATIVE-ION FORMATION POSITIVE-ION-SURFACE SCATTERING TWO-ELECTRON TRANSFER TIME-DEPENDENT ANDERSON-NEWNS MODEL CORRELATION ENERGY H <sup>+</sup> + CESIATED W(100) GOOD AGREEMENT WITH EXPERIMENT		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A theoretical investigation of negative-ion formation from positive-ion-surface scattering is presented from a unified point of view. Based on the time-dependent Anderson-Newns model, the correlation energy U is seen to play an important role in the two-electron transfer process. Calculations of the probability of negative-ion formation are in good agreement with experiments on the conversion of H <sup>+</sup> (D <sup>+</sup> ) to H <sup>-</sup> (D <sup>-</sup> ) by scattering from a cesiated W(100) surface.		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Physics Letters A, in press

NEGATIVE-ION FORMATION FROM SURFACE SCATTERING  
AND THE ANDERSON CORRELATION ENERGY U

Kai-Shue Lam\*, K. C. Liu and Thomas F. George  
Department of Chemistry  
University of Rochester  
Rochester, New York 14627

ICSU Classification #: 68, 78, 79



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

\*Present address: Department of Physics  
California Polytechnic State University  
3801 West Temple Avenue  
Pomona, California 91768

### Abstract

A theoretical investigation of negative-ion formation from positive-ion-surface scattering is presented from a unified point of view. Based on the time-dependent Anderson-Newns model, the correlation energy  $U$  is seen to play an important role in the two-electron transfer process. Calculations of the probability of negative-ion formation are in good agreement with experiments on the conversion of  $H^+(D^+)$  to  $H^-(D^-)$  by scattering from a cesiated W(100) surface.

Charge-exchange processes arising from monoenergetic ion beams scattered from solid surfaces have recently been the subject of much experimental<sup>1-5</sup> and theoretical<sup>6-10</sup> interest. Most of these studies focus on the process of ion neutralization, involving the transfer of a single electron from the surface to the singly-charged ion.<sup>1-3,6-10</sup> However, owing to the increase of applications involving the transfer of two electrons in a variety of situations, such as plasma fusion<sup>11,12</sup> and the generation of high-energy neutral beams,<sup>11</sup> attention has also been directed to processes of negative-ion formation.<sup>4,5,13</sup>

For ion neutralization, the majority of theoretical work has been based on the Anderson-Newns model,<sup>14</sup> originally proposed for the explanation of localized magnetization in transition metal alloys and subsequently applied to various other problems such as chemisorption on metals<sup>15</sup> and mixed valence in solids.<sup>16,17</sup> In these applications of this model, the correlation energy  $U$ , arising from the Coulomb repulsion between the two electrons of opposite spin in the same discrete level, plays a crucial role. However, in studies of ion neutralization (or the reverse process of atom ionization), this important quantity is either implicitly assumed to be infinite<sup>6</sup> or completely ignored.<sup>7-9</sup> Such approaches are thus incapable of accounting for negative ion formation:  $U = \infty$  completely suppresses the transfer of a second electron, while if only single-electron transfer is considered,  $U$  is irrelevant.

Previous theoretical work<sup>13</sup> on negative-ion formation has centered on the time-dependent width and shift of the valence level of the impact ion, where the time dependence is due to the motion of the ion. It was assumed that first neutralization occurs via a nonresonant Auger process, while the

nuclear motion of the ion can subsequently bring the valence level into resonance with a band state and thus effect the transfer of the second electron. The quantity  $U$  again does not play an important role in this theory.

In the present work, we use the Anderson-Newns model to examine the effects of  $U$  on the two-electron transfer process in relation to the dynamics of the nuclear motion of the impact ion. This represents the first time that charge-exchange processes in surface scattering have been considered from a unified point of view. We shall see that the negative-ion formation depends crucially on the finite value of  $U$ , and in fact, recent experimental results<sup>5</sup> can be explained in terms of our theory.

The time-dependent Anderson-Newns Hamiltonian including the correlation energy term is given as

$$H(t) = H_0 + V(t), \quad (1)$$

where

$$H_0 = \sum_{\sigma} \epsilon_d c_{d\sigma}^{\dagger} c_{d\sigma} + \sum_{k\sigma} \epsilon_k c_{k\sigma}^{\dagger} c_{k\sigma} + U n_{d\sigma} n_{d,-\sigma} \quad (2)$$

$$V(t) = \sum_{k\sigma} [V_{kd}(t) c_{d\sigma}^{\dagger} c_{k\sigma} + V_{kd}^*(t) c_{k\sigma}^{\dagger} c_{d\sigma}]. \quad (3)$$

The indices  $d$  and  $k$  denote the valence state of the impact ion and the conduction band states of the solid, respectively,  $\sigma$  is the spin index and  $n_{d\sigma} = c_{d\sigma}^{\dagger} c_{d\sigma}$ . The interaction Hamiltonian,  $V(t)$ , responsible for the electron transfer between the band and valence states, is the only explicitly time-dependent part. The motion of the projectile ion can be phenomenologically



taken into account by using the specific form<sup>6</sup>

$$V_{kd}(t) = V_{kd} e^{-\lambda |t|}, \quad (4)$$

where  $\lambda$ , the sole dynamical parameter in our model, is directly proportional to the normal impact velocity. Hence  $\lambda$  controls the duration of the bound-continuum interaction, while  $V_{kd}$  (time-independent) determines its strength.

To lowest order, the perturbative solution for the time-evolution operator  $\tilde{T}$  in the interaction picture, which contributes to the S-matrix for negative-ion formation, is given as

$$\tilde{T}^{(2)}(t, -\infty) = (-i)^2 \int_{-\infty}^t dt' \tilde{V}(t') \int_{-\infty}^t dt'' \tilde{V}(t''), \quad (5)$$

where

$$\tilde{V}(t) = e^{iH_0 t} \hat{V}(t) e^{-iH_0 t} \quad (6)$$

is  $V(t)$  expressed in the interaction picture. At  $t = -\infty$ , the ion is taken to be infinitely far from the surface, and  $t = 0$  is the instant of impact. It is apparent from Eq. (3) that, since negative-ion formation involves ultimately the transfer of two electrons, only terms of even powers in  $\hat{V}$  in the expansion of  $\tilde{T}$  contribute to the S-matrix.

Figure 1 illustrates schematically the electron-transfer process to lowest, i.e., second, order as contained in Eq. (5). Physically, every factor of  $\tilde{V}(t)$  corresponds to the transfer of one electron from the band level  $\epsilon_k$  to the  $\epsilon_d$  level of the ion, or the conjugate process of electron transfer from  $\epsilon_d$  to  $\epsilon_k$ . Therefore, the contribution to the S-matrix from

higher-order terms of  $\tilde{T}$  can be diagrammatically generated in a straightforward manner. Such terms (neglected in this work), however, may involve those final states in which the solid becomes electronically excited, with levels higher than the Fermi level occupied. In general, low-ordered processes are favored by high impact velocities (large  $\lambda$ ).

To lowest order, the time-dependent probability for negative-ion formation is then given as

$$P(t) = \int_{\epsilon_L}^{\epsilon_F} d\epsilon \int_{\epsilon_L}^{\epsilon_F} d\epsilon' \rho(\epsilon) \rho(\epsilon') |\langle k_+ k'_+ | \tilde{T}^{(2)}(t, -\infty) | I \rangle|^2, \quad (7)$$

where  $\epsilon_F$  and  $\epsilon_L$  are the Fermi energy and conduction band edge of the solid, respectively,  $\rho(\epsilon)$  is the density of states of the band, and the initial and final states,  $|I\rangle$  and  $|k_+ k'_+\rangle$  respectively, are described in Fig. 1. The matrix element in Eq. (7) can be evaluated by using Eqs. (2)-(6) to give the result

$$\langle k_+ k'_+ | \tilde{T}^{(2)}(t, -\infty) | I \rangle = -\frac{v^2}{2} \frac{1}{\lambda + i(U - \epsilon - \epsilon')/2} e^{[2\lambda + i(U - \epsilon - \epsilon')]t} \left( \frac{1}{\lambda - i\epsilon} + \frac{1}{\lambda - i\epsilon'} \right), \quad t \leq 0 \quad (8a)$$

$$\begin{aligned} &= -\frac{v^2}{2} \frac{1}{\lambda + i(U - \epsilon - \epsilon')/2} \left( \frac{1}{\lambda - i\epsilon} + \frac{1}{\lambda - i\epsilon'} \right) - \frac{v^2}{2} \frac{1}{\lambda - i(U - \epsilon - \epsilon')/2} \\ &\times \left( \frac{1}{\lambda + i\epsilon} + \frac{1}{\lambda + i\epsilon'} \right) \left( e^{-[2\lambda + i(U - \epsilon - \epsilon')]t} - 1 \right) \\ &+ 2v^2 \lambda \left\{ \frac{1}{(\lambda^2 + \epsilon^2)(\lambda - i(U - \epsilon))} \left( e^{-[\lambda - i(U - \epsilon)]t} - 1 \right) \right. \\ &\left. + \frac{1}{(\lambda^2 + \epsilon'^2)(\lambda - i(U - \epsilon'))} \left( e^{-[\lambda - i(U - \epsilon')]t} - 1 \right) \right\} \quad t > 0 \quad (8b) \end{aligned}$$

where  $\epsilon = \epsilon_k - \epsilon_d$  and  $\epsilon' = \epsilon_{k'} - \epsilon_d$  with  $\epsilon_d$  set as the zero of energy, and  $v = v_{kd}$  is assumed to be independent of energy. As a first approximation, we have also assumed that  $U$  and  $\epsilon_d$  are constant within the collision region. This latter assumption is not expected to affect the results qualitatively.

We now turn to calculations of the probability  $P(t)$  of negative-ion formation and its limiting value at  $t \rightarrow \infty$ ,  $P(\infty)$ , for various choices of the parameters  $U$  and  $\lambda$ .  $P(\infty)$  represents the experimental observation of outgoing negative ions after the scattering event is complete, and  $P(t)$  reflects the behavior of the transient states. The following fixed numerical values were adopted:  $\epsilon_L = -10$  eV,  $\epsilon_F = -0.06$  eV and  $\epsilon = \epsilon_L^2 = 0.43$  eV, where we have assumed the band to have no important structure so that the density of states  $\rho$  can be taken as constant in the integration over the band in Eq. (7). Although the last two numbers have been used previously in a study of charge transfer in the Na/W(100) system,<sup>8</sup> they have no special significance in our present work, which is to investigate in general the effects of the variation of  $\lambda$  and  $U$ , especially the latter.

Intuitively, we expect the probability  $P(\infty)$  to be small when the repulsive correlation energy  $U$  is large. However,  $P(\infty)$  also depends on  $\lambda$ , i.e., on the velocity of the impact ion. As shown in Fig. 2, there is a peak for  $P(\infty)$  at a small value of  $\lambda$  for each  $U$ . The explanation for this is that since small  $\lambda$  implies a long duration of interaction whereby from the uncertainty principle the resonance requirement is stringent, it is impossible for the second electron in the solid to overcome the barrier  $U$ . On the other hand, large  $\lambda$  can ease the resonance requirement -- energy conservation can be violated for short-duration processes -- but it also limits the actual time available for electron transfer, resulting in a small  $P(\infty)$ . As a consequence, for each  $U$  there exists an optimal value of

$\lambda$ ,  $\lambda_m$ , for which the probability attains a maximum. Moreover,  $\lambda_m$  increases as  $U$  is increased, due to the fact that a shorter interaction time is favorable for the second electron to transfer non-resonantly as  $U$  becomes larger.

The close relationship between resonant electron transfer and the ion velocity becomes obvious when we look at the time evolution of the probability of negative-ion formation. Figure 3 displays  $P(t)$  for various values of  $U$  with fixed  $\lambda$ . Each probability curve has a peak at very short time  $t_m$  ( $\sim 0.2$  femtoseconds). We see that  $t_m$  is smaller for larger  $U$ , in accordance with the arguments given above. In addition, the degree of transient negative-ion formation, measured by the ratio  $P(t_m)/P(\infty)$ , is more pronounced for larger  $U$ . Figure 4 provides yet another manifestation of the striking transient behavior; namely, although there is an optimal  $\lambda_m$  at each  $U$  for the limiting value of the probability,  $P(\infty)$ , this is not necessarily the case in the transient region.

To test our theory, we shall compare it with measurements on the conversion of  $H^+(D^+)$  to  $H^-(D^-)$  by scattering from a cesiated W(100) surface at different grazing angles  $\theta$ .<sup>5</sup> For this purpose, we identify  $\lambda$  as  $v \cos \theta$ , where  $v$  is the magnitude of the velocity, and phenomenologically introduce a velocity-dependent interaction,  $\Delta = 0.43 \exp(-0.01v)$  (in the unit of eV), to account for the loss of particles due to penetration into the surface. The variation of  $\theta$  is thus equivalent to the variation of the normal impact velocity  $\lambda$ . Our results, given in Fig. 5, are in qualitative agreement with the experimental ones,<sup>5</sup> where for all incident energies the conversion probability goes

through a maximum. Quantitative comparisons have not been attempted since precise information on critical parameters, especially  $\Delta$ , is still lacking. For the cesiated W(100) surface, among other complications leading to unreliable data for parameters is the theoretical evidence of a lowering of the work function by multiple dipole formation.<sup>18,19</sup>

In this work we have demonstrated, through varying the dynamical conditions of the impact ion, the significance of the correlation energy  $U$  in negative-ion formation from positive-ion-surface scattering. Though  $U$  in general decreases the probability for negative-ion formation, one can always exploit the experimentally controllable dynamical conditions (varying  $v$  and  $\theta$ ) to achieve an optimal result for a given system. Moreover, there may even be the possibility of exploiting the characteristic transient behavior of  $P(t)$ , since for finite  $U$ ,  $P(t_m)$  is always larger than  $(\infty)$  except for very large  $\lambda$ . For very small  $\lambda$ , on the other hand, our perturbation approach may not yield correct results, since the long interaction times then allowed may require higher-order processes than the second-order one be considered. Our results have been shown to be in good qualitative agreement with experiments. A more elaborate calculation is needed which takes into account the lowering of the valence level of the ion near the surface<sup>18,19</sup> is needed for quantitative comparison with experiments.

#### Acknowledgments

This research was supported in part by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Grant AFOSR-82-0046, and the Office of Naval Research. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. TFG acknowledges the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Award (1975-84) and the John Simon Guggenheim Foundation for a Fellowship (1983-84).

### References

1. H. H. Brogersma and T. M. Buck, Surf. Sci. 53, 649 (1975).
2. R. L. Erickson and D. P. Smith, Phys. Rev. Lett. 34, 297 (1975).
3. E. G. Overbosch, B. Rasser, A. D. Tenner and J. Los, Surf. Sci. 92, 310 (1980).
4. E. G. Overbosch and J. Los, Surf. Sci. 108, 117 (1981).
5. J. N. M. van Wunnik, B. Rasser and J. Los, Phys. Lett. 87A, 288 (1982); J. Los, E. A. Overbosch and J. N. M. van Wunnik in Proceedings of the Second International Symposium on the Production and Neutralization of Negative Hydrogen Ions and Beams (Brookhaven, 1980).
6. W. Bloss and D. Hone, Surf. Sci. 72, 277 (1978).
7. Y. Muda and T. Hanawa, Surf. Sci. 97, 283 (1980).
8. R. Brako and D. M. Newns, Surf. Sci. 108, 253 (1981); Vacuum 32, 39 (1982).
9. K. L. Sebastian, V. C. Jyothi Bhasu and T. B. Grimley, Surf. Sci. 110, L571 (1981).
10. J. C. Tully, Phys. Rev. B 16, 4324 (1977).
11. K. H. Berkner, R. V. Pyle and J. W. Stearns, Nucl. Fusion 15, 249 (1979).
12. K. Wiesemann, K. Prelec and Th. Sluyters, J. Appl. Phys. 48, 2668 (1977).
13. B. Rasser, J. N. M. van Wunnik and J. Los, Surf. Sci. 118, 697 (1982).
14. P. W. Anderson, Phys. Rev. 124, 41 (1961).
15. D. M. Edwards and D. M. Newns, Phys. Lett. 24A, 236 (1967); D. M. Newns, Phys. Rev. 178, 1123 (1969).
16. B. Coqblin and J. R. Schrieffer, Phys. Rev. 185, 847 (1969).
17. J. H. Jefferson and K. W. Stevens, J. Phys. C 9, 2151 (1976).
18. K. H. Kingdon and I. Langmuir, Phys. Rev. 21, 380 (1923).
19. E. Wimmer, A. J. Freeman, N. Weinert, H. Krakauer, J. R. Hiskes and A. M. Karo, Phys. Rev. Lett. 48, 1128 (1982).

### Figure Captions

Fig. 1. Schematic diagram of electron transfer in negative-ion formation.

$\epsilon_U$  is the upper and  $\epsilon_L$  the lower edge of the band. (a) Initial state  $|I\rangle$ : valence state  $\epsilon_d$  empty, band filled up to the Fermi level  $\epsilon_F$ . (b) Intermediate states  $|k_\uparrow\rangle$  and  $|k'_\downarrow\rangle$ : states corresponding to the neutralized atom; one electron transferred from the  $\epsilon_k$  or  $\epsilon_{k'}$  level to the  $\epsilon_d$  level. The arrows denote the spin states of the electrons, and the solid and hollow circles represent electrons and holes, respectively. (c) Final states  $|k_\uparrow k'_\downarrow\rangle$ : negative-ion states; two electrons transferred to the  $\epsilon_d$  level.

Fig. 2.  $P(\infty)$  vs  $\lambda$  for various values of  $U$ . Energy is in the unit of eV.

Fig. 3.  $P(t)$  vs  $t$  for various values of  $U$  with fixed  $\lambda$ . As  $U$  increases, the characteristic short-time behavior becomes more pronounced. Energy is in the unit of eV, and time is in the unit of  $6.59 \times 10^{-16} \text{ sec } (\frac{\hbar}{\text{eV}})$

Fig. 4.  $P(t)$  vs  $t$  for various values of  $\lambda$  with fixed  $U$ . The units are the same as in Fig. 3.

Fig. 5.  $P(\infty)$  vs  $\theta$ , the incident angle of impact.  $v_1 < v_2 < v_3 < v_4$ .




FIG. 1






FIG. 2




FIG. 3




FIG. 4




FIG. 5

DL/413/83/01  
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1
Dr. David L. Nelson Chemistry Program Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1		

DL/413/83/01  
056/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Surface Chemistry Division (6170)  
455 Overlook Avenue, S.W.  
Washington, D.C. 20375

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. Chia-wei Woo  
Department of Physics  
Northwestern University  
Evanston, Illinois 60201

Dr. Robert M. Hexter  
Department of Chemistry  
University of Minnesota  
Minneapolis, Minnesota

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. Adolph B. Amster  
Chemistry Division  
Naval Weapons Center  
China Lake, California 93555

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60201

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. S. L. Bernasek  
Princeton University  
Department of Chemistry  
Princeton, New Jersey 08544

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. F. Carter  
Code 6132  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Richard Colton  
Code 6112  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Professor R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York ~~22217~~  
12301

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 5570  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
California Institute of Technology  
Division of Chemistry and Chemical  
Engineering  
Pasadena, California 91125

Professor E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
Southampton University  
Southampton SO9 5NH  
Hampshire, England

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

Dr. Richard Smardzewski  
Code 6130  
Naval Research Laboratory  
Washington, D.C. 20375

DL/413/83/01  
056/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. P. Hansma  
Physics Department  
University of California  
Santa Barbara, California 93106

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Professor T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Professor Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Captain Lee Myers  
AFOSR/NC  
Bolling AFB  
Washington, D.C. 20332

Professor Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Professor D. Hercules  
University Pittsburgh  
Chemistry Department  
Pittsburgh, Pennsylvania 15260

Professor N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. G. D. Stein  
Mechanical Engineering Department  
Northwestern University  
Evanston, Illinois 60201

Professor A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Professor G. H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. David Squire  
Army Research Office  
P.O. Box 12211  
Research Triangle Park, NC 27709



ATE  
LMED  
8